

|                                                                                                                                                                                                                                                                                                                 |       |                                                                                                                                                                                                                                                                 |       |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <b>Structural Biology of Riboswitch-mediated Gene Regulation and Argonaute-mediated Gene Silencing</b><br>D. J. Patel                                                                                                                                                                                           | 1     | <b>Building Biologically Active Nucleic Acid Nanocomplexes</b><br>C. I. E. Smith, K. E. Lundin, O. E. Simonson, P. M. D. Moreno, M. G. Svahn, M. Wenska and R. Strömberg                                                                                        | 27–28 |
| <b>Characterization and Use of Tricyclic Fluorescent Nucleic Acid Base Analogues</b><br>K. Börjesson, P. Sandin, B. Albinsson, J. Mårtensson, T. Brown and L. M. Wilhelmsson                                                                                                                                    | 3–4   | <b>Recognizing and Controlling Biomolecules with “Smart” Hybridization-based Switches</b><br>O. Seitz, T. N. Grossmann, S. Thurley and L. Röglin                                                                                                                | 29–30 |
| <b>Nucleic acid duplexes with zippers of additional nucleobases and aromatics in minor or major groove</b><br>P. Nielsen, K. I. Shaikh, L. J. Nielsen, N. K. Andersen, L. Spáčilová, H. Nielsen, P. Kočalka and M. S. Christensen                                                                               | 5–6   | <b>PNA-peptide conjugates as intracellular gene control agents</b><br>G. D. Ivanova, M. M. Fabani, A. A. Arzumanov, R. Abes, H. Yin, B. Lebleu, M. Wood and M. J. Gait                                                                                          | 31–32 |
| <b>DNA and RNA Quadruplex ligands</b><br>A. D. Cian, J. Gros, A. Guédin, M. Haddi, S. Lyonnais, L. Guittat, J.-F. Riou, C. Trentesaux, B. Saccà, L. Lacroix, P. Alberti and J.-L. Mergny                                                                                                                        | 7–8   | <b>Structural and functional prerequisites for ribosomal nascent peptide acceptors: Attempts to decipher the nature of the ribosome’s catalysis of peptide bond formation</b><br>B. Y. Michel, K. S. Krishnakumar, M. Johansson, M. Ehrenberg and P. Strazewski | 33–34 |
| <b>G-quadruplex forming oligonucleotides as finely tunable aptamers: towards better DNA mimics</b><br>D. Montesarchio                                                                                                                                                                                           | 9–10  | <b>The acetal levulinyl ester (ALE) group for the 2'-hydroxyl protection of ribonucleosides and the synthesis of oligoribonucleotides</b><br>J. G. Lackey and M. J. Damha                                                                                       | 35–36 |
| <b>The use of light to investigate and modulate DNA and RNA conformations</b><br>D. Sen                                                                                                                                                                                                                         | 11–12 | <b>Triplex glue by synthesizing conjugated flexible intercalators</b><br>E. B. Pedersen, A. M. A. Osman, D. Globisch, M. Paramasivam, S. Cogoi, N. Bomholt, P. T. Jørgensen, L. E. Xodo and V. V. Filichev                                                      | 37–38 |
| <b>Manipulation of the peptide-binding specificity of an RNA in a rational manner by combinations of specificity-altering mutations</b><br>K. Harada, M. Sugaya, F. Nishimura and A. Katoh                                                                                                                      | 13–14 | <b>New strategies for the synthesis of unmodified and modified oligonucleotides</b><br>M. Sekine, A. Ohkubo and K. Seio                                                                                                                                         | 39–40 |
| <b>Quantitative, Sensitive Analysis of DNA and RNA</b><br>C. R. Cantor                                                                                                                                                                                                                                          | 15    | <b>RNA interference in silencing of genes of Alzheimer’s disease in cellular and rat brain models</b><br>M. Sierant, K. Kubiak, J. Kazmierczak-Baranska, A. Patuszynska, T. Kuwabara, M. Warashina, B. Nacmias, S. Sorbi and B. Nawrot                          | 41–42 |
| <b>Artificial molecular switches made from DNA</b><br>E. Friedrichs, A. Tsokou, R. Jungmann and F. Simmel                                                                                                                                                                                                       | 17–18 | <b>Synthesis of the new nucleoside analogue connecting 2-amino-6-vinylpurine to the 2'-deoxyribose skeleton via the methylene linker</b><br>Y. Kurose, Y. Taniguchi, F. Nagatsugi and S. Sasaki                                                                 | 43–44 |
| <b>Construction of a photo-switchable gene for turning on and off gene expression with light irradiation</b><br>X. Liang, K. Fujioka, Y. Tsuda, R. Wakuda and H. Asanuma                                                                                                                                        | 19–20 | <b>Potent single stranded RNA inhibition</b><br>T. Koch                                                                                                                                                                                                         | 45    |
| <b>DNA Controlled Assembly of Liposomes</b><br>U. Jakobsen, A. C. Simonsen and S. Vogel                                                                                                                                                                                                                         | 21–22 | <b>Click chemistry and Oligonucleotides: How a simple reaction can do so much</b><br>F. Morvan, A. Meyer, G. Pourceau, S. Vidal, Y. Chevlot, E. Souteyrand and J.-J. Vasseur                                                                                    | 47–48 |
| <b>Zinc finger protein-based detection system of PCR products for pathogen diagnosis</b><br>Y. Osawa, H. Motoki, T. Matsuo, M. Horiuchi, K. Sode and K. Ikebukuro                                                                                                                                               | 23–24 | <b>Endogenous cytosine methylation and the formation of carcinogen–DNA adducts</b><br>N. Tretyakova, R. Guza and B. Matter                                                                                                                                      | 49–50 |
| <b>LNA<sup>®</sup> incorporated siRNAs exhibit lower off-target effects compared to 2'-OMethoxy in Cell Phenotypic Assays and Microarray Analysis</b><br>N. Puri, X. Wang, R. Varma, C. Burnett, L. Beauchamp, D. M. Batten, M. Young, V. Sule, K. Latham, T. Sendera, C. Echeverri, C. Sachse and S. Magdaleno | 25–26 | <b>A Whole Base-Labile Strategy for RNA Synthesis with 2'-O-acetalester Protections</b><br>T. Lavergne, A. Martin, F. Debart and J.-J. Vasseur                                                                                                                  | 51–52 |

|                                                                                                                                                                                                                                                                          |       |                                                                                                                                                                                                                                                                                                       |         |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Novel base-functionalized DNA. Efficient methodology for construction and bioanalytical applications</b><br>M. Hocek, M. Vrábel, H. Cahová, J. Riedl, L. Kalachová, H. Pivoňková, P. Horáková, L. Havran and M. Fojta                                                 | 53–54 | <b>Synthesis and properties of (<math>\alpha</math>-<i>P</i>-borano)-nucleoside 5'-triphosphate analogues as potential antiviral agents</b><br>M. A. Cheek, M. I. Dobrikov, C. K. Wennefors, Z. Xu, S. N. Hashmi, X. Shen and B. R. Shaw                                                              | 81–82   |
| <b>Recent Developments in the Synthesis of RNA Oligonucleotides for Potential Therapeutic Applications</b><br>J. Cieślak, A. Grajkowski, C. Ausin and S. L. Beaucage                                                                                                     | 55–56 | <b>Nucleoside Diphosphate Prodrugs</b><br>C. Meier, H. J. Jessen and J. Balzarini                                                                                                                                                                                                                     | 83–84   |
| <b>Repair of DNA-protein crosslink damage: Coordinated actions of nucleotide excision repair and homologous recombination</b><br>H. Ide, T. Nakano, A. M. H. Salem, H. Terato, S. P. Pack and K. Makino                                                                  | 57–58 | <b>Inhibition of Orotidine-5'-monophosphate decarboxylase – Discoveries and lessons</b><br>L. P. Kotra and E. F. Pai                                                                                                                                                                                  | 85–86   |
| <b>Characterization of endogenous human Argonautes and their miRNA partners in RNA silencing</b><br>M. C. Siomi and H. Siomi                                                                                                                                             | 59–60 | <b>Multifunctional envelope type nano device for non-viral gene delivery: Concept and application for nanomedicine</b><br>H. Harashima                                                                                                                                                                | 87      |
| <b>Physicochemical Stability of NOX-E36, a 40mer L-RNA (Spiegelmer) for Therapeutic Applications</b><br>C. Maasch, K. Buchner, D. Eulberg, S. Vonhoff and S. Klussmann                                                                                                   | 61–62 | <b>Preparation of Mannosylated Oligoribonucleotides</b><br>Y. Zhao and H. (Tony) Yan                                                                                                                                                                                                                  | 89–90   |
| <b>Inhibition of picornaviruses by means of RNA interference</b><br>D. Rothe, D. Werk, M. Dutkiewicz, S. Schubert, H.-P. Grunert, H. Zeichhardt, V. A. Erdmann, H. Fehner and J. Kurreck                                                                                 | 63–64 | <b>Tyrosine-modified PEI: A novel and highly efficient vector for siRNA delivery in mammalian cells</b><br>G. Creusat and G. Zuber                                                                                                                                                                    | 91–92   |
| <b>Molecular engineering of DNA bases: Building block for functional molecular probes in biomedicine</b><br>Y. Wu, Y. Kim and W. Tan                                                                                                                                     | 65–66 | <b>Designer DNA Architectures for Nanobiotechnology</b><br>Y. Liu and H. Yan                                                                                                                                                                                                                          | 93–94   |
| <b>Protein-Facilitated Ribozyme Folding and Catalysis</b><br>N. Zingler, A. Solem and A. M. Pyle                                                                                                                                                                         | 67–68 | <b>DNA electron transfer mechanism and dynamics</b><br>F. D. Lewis                                                                                                                                                                                                                                    | 95–96   |
| <b>2'-O-Methoxyethyl/2'-Fluoro Modified Oligonucleotides Result in More Potent Inhibition of micro RNA-122 in Vivo: A Target implicated in HCV Replication</b><br>B. Bhat, C. Esau, S. Davis, S. Propp, G. Kinberger, H. Gaus, S. Freier, E. E. Swayze and C. F. Bennett | 69    | <b>Addressable Molecular Node Assembly – Functional DNA Nanostructures</b><br>J. Tumpane, E. P. Lundberg, L. M. Wilhelmsson, T. Brown and B. Nordén                                                                                                                                                   | 97–98   |
| <b>An efficient fluorescent method for selective detection of mature miRNA species</b><br>Y. Kato                                                                                                                                                                        | 71–72 | <b>Synthesis of 2',3'-Cyclohexene Bicyclic Nucleoside Analogues as Antiviral Compounds</b><br>A. Díaz-Rodríguez, S. Fernández, Y. S. Sanghvi, E. A. Theodorakis, V. Gotor and M. Ferrero                                                                                                              | 99–100  |
| <b>In vitro selection of a DNazyme with three modified nucleotides</b><br>M. Hollenstein, C. Hipolito, C. Lam and D. Perrin                                                                                                                                              | 73–74 | <b>Preparation of Nucleoside-Carbohydrate Phosphodiester Prodrug Analogues by Chemoenzymatic Procedure</b><br>T. Rodríguez-Pérez, S. Fernández, Y. S. Sanghvi, V. Gotor and M. Ferrero                                                                                                                | 101–102 |
| <b>Transition state analogues in quorum sensing and SAM recycling</b><br>V. L. Schramm, J. A. Gutierrez, G. Cordovano, I. Basu, C. Guha, T. J. Belbin, G. B. Evans, P. C. Tyler and R. H. Furneaux                                                                       | 75–76 | <b>RNA Synthesis: Phosphoramidites for RNA synthesis in the reverse direction. Highly efficient synthesis and application to convenient introduction of ligands, chromophores and modifications of synthetic RNA at the 3'- end</b><br>S. C. Srivastava, D. Pandey, N. P. Srivastava and S. P. Bajpai | 103–104 |
| <b>Chemistry and Structure-Activity Relationship of Antibacterial Nucleoside Natural Products</b><br>S. Ichikawa and A. Matsuda                                                                                                                                          | 77–78 | <b>Studies of sequence-specific recognition and interaction of bishairpin polyamide minor groove binders with target DNA duplexes</b><br>A. S. Boutorine, W. Buchmann, L. Halby and E. De Pauw                                                                                                        | 105–106 |
| <b>Development of A<sub>3</sub> Adenosine Receptor Ligands</b><br>L. S. Jeong                                                                                                                                                                                            | 79–80 | <b>Binding of two bis-bipyridine minor groove binders to a DNA template in the presence of Cu<sup>2+</sup> ions</b><br>J. A. Brazier, I. Onishi and S. Sasaki                                                                                                                                         | 107–108 |

|                                                                                                                                                                                                                              |         |                                                                                                                                                                                                                                                            |         |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Recognition of Homopyrimidine Mismatches by Distance-Constrained Macrocyclic bisintercalators</b><br>M. Bahr, V. Gabélica, A. Granzhan, E. Weinhold and M.-P. Teulade-Fichou                                              | 109–110 | <b>Formation of a stable triplex incorporating a CG interrupting site by a new WNA derivative containing 3-aminopyrazole as a nucleobase</b><br>Y. Uchida, Y. Taniguchi, E. Aoki, M. Togo and S. Sasaki                                                    | 137–138 |
| <b>The interaction between the purine motif triplex and the triplex DNA-binding domain of <i>Saccharomyces cerevisiae</i> Stm1 protein</b><br>T. Katayama and H. Torigoe                                                     | 111–112 | <b>Strand invasion properties and serum stability of <math>\alpha</math>-tricyclo-DNA</b><br>S. P. Scheidegger and C. J. Leumann                                                                                                                           | 139–140 |
| <b>Effect of cationic comb-type copolymer on the B- Z transition of poly(dG-dC)-poly(dG-dC)</b><br>N. Shimada, A. Kano and A. Maruyama                                                                                       | 113–114 | <b>Single-strand binding protein enhances invasion of a PNA strand to double-stranded DNA</b><br>T. Ishizuka and M. Komiyama                                                                                                                               | 141–142 |
| <b>A pteridine derivative with electron-withdrawing groups for binding and sensing of nucleobases in AP site-containing DNA duplexes</b><br>E. Kanai, S. Nishizawa and N. Teramae                                            | 115–116 | <b>Introduction of linkers into PNA for versatile applications</b><br>Y. Aiba and M. Komiyama                                                                                                                                                              | 143–144 |
| <b>Effect of an alkyl amino group on the binding of 1,8-naphthyridines to AP site-containing DNA duplexes</b><br>T. Ichihashi, Y. Sato, T. Seino, S. Nishizawa and N. Teramae                                                | 117–118 | <b><i>Iso</i>-Thioacetamido nucleic acids (<i>iso</i>TANA): Synthesis and DNA binding studies of Lyxo/Ribo oligonucleotides</b><br>S. Gokhale and V. A. Kumar                                                                                              | 145–146 |
| <b>Competitive binding of small ligands to nucleobases in AP site-containing DNA duplexes</b><br>T. Kageyama, Y. Sato, S. Nishizawa and N. Teramae                                                                           | 119–120 | <b>Site-specific insertion of nitroxide-spin labels into DNA probes by click chemistry for structural analyses by ELDOR spectroscopy</b><br>M. Flaender, G. Sicoli, Th. Fontecave, G. Mathis, C. Saint-Pierre, Y. Boulard, S. Gambarelli and D. Gasparutto | 147–148 |
| <b>Development of label-free molecular beacons based on abasic site-binding fluorescence molecules</b><br>Y. Sato, S. Nishizawa and N. Teramae                                                                               | 121–122 | <b>A click chemistry approach towards nucleic acid major groove functionalization</b><br>N. K. Andersen, L. Spáčilová, M. D. Jensen, P. Kočalka, F. Jensen and P. Nielsen                                                                                  | 149–150 |
| <b>A surface plasmon resonance sensor based on 3,5-diaminopyrazine with a high selectivity for thymine in AP site-containing DNA duplex</b><br>S. Miura, K. Ono, M. Watanabe, S. Nishizawa and N. Teramae                    | 123–124 | <b>Dynamics of 2-<i>N</i>-<i>tert</i>-Butylaminoxyladenosine Incorporated into Oligodeoxynucleotides</b><br>M. Aso, M. Kurita, J. W. Mirc, T. Kaneko, Y. Higuchi, N. Koga and H. Suemune                                                                   | 151–152 |
| <b>Mismatch recognition in PNA double-duplex invasion</b><br>Y. Miyajima, T. Ishizuka and M. Komiyama                                                                                                                        | 125–126 | <b>Nitroxide Spin labeled RNA for long range distance measurements by EPR-PELDOR</b><br>O. Frolow, B. Endeward, O. Schiemann, T. F. Prisner and J. W. Engels                                                                                               | 153–154 |
| <b>Crystal structures of DNA duplexes stabilized by bicyclic-C residues</b><br>T. Haraguchi, S. Shimizu, X. Ma, T. Kurose, E. C. M. Juan, A. Ohkubo, M. Sekine, T. Shibata, C. L. Millington, D. M. Williams and A. Takénaka | 127–128 | <b>New triarylamine-based far-red DNA stainers with high two-photon absorption properties</b><br>G. Bordeau, E. Faurel, C. Allain, R. Lartia, F. Schmidt, C. Fiorini-Debuisschert, F. Charra, G. Metge, P. Tauc and M.-P. Teulade-Fichou                   | 155–156 |
| <b>Self-Avoiding molecular Recognition Systems (SAMRS)</b><br>S. Hoshika, F. Chen, N. A. Leal and S. A. Benner                                                                                                               | 129–130 | <b>Role of Molecular Crowding in perturbing Quadruplex-Watson Crick Duplex Equilibrium</b><br>N. Kumar and S. Maiti                                                                                                                                        | 157–158 |
| <b>Finding at-DNA – Kinetic Recognition of Long Adenine-Thymine Stretches by Metal-Ligand Complexes</b><br>P. Nordell, F. Westerlund, A. Reymer, B. Nordén and P. Lincoln                                                    | 131–132 | <b>Protein hnRNPA1 binds to a critical G-rich element of <i>KRAS</i> and unwinds G-quadruplex structures: implications in transcription</b><br>L. Xodo, M. Paramasivam, A. Membrino and S. Cogo                                                            | 159–160 |
| <b>Unlocked nucleic acid (UNA) and UNA derivatives: Thermal denaturation studies</b><br>T. B. Jensen, N. Langkjær and J. Wengel                                                                                              | 133–134 | <b>Structural stability analysis of the intermediates in the folding pathway of human telomeric hybrid-1 G-quadruplex based on fragment molecular orbital method</b><br>H. Yagi, T. Mashimo, Y. Sannohe and H. Sugiyama                                    | 161–162 |
| <b>Synthesis of 3'- and 6'-functionalised Locked Nucleic Acid (LNA) analogues</b><br>P. K. Sharma, G. Enderlin, S. Kumar and P. Nielsen                                                                                      | 135–136 |                                                                                                                                                                                                                                                            |         |

|                                                                                                                                                                                                                                          |         |                                                                                                                                                                                                                                  |         |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>New platinum(II) complexes targeting the loops of the human telomeric G-quadruplex</b><br>H. Bertrand, S. Bombard, D. Monchaud and M.-P. Teulade-Fichou                                                                               | 163–164 | <b>Generation of plants resistant to tomato yellow leaf curl virus by using artificial zinc-finger proteins</b><br>Y. Koshino-Kimura, K. Takenaka, F. Domoto, Y. Aoyama and T. Sera                                              | 189–190 |
| <b>Ligand binding to tetra-end-linked (TGGGGT)<sub>4</sub> G-quadruplexes: an electrospray mass spectroscopy study</b><br>J. Amato, G. Oliviero, N. Borbone, S. D'Errico, G. Piccialli, P. Mailliet, F. Rosu, E. D. Pauw and V. Gabelica | 165–166 | <b>Synthesis of locked S-type and locked N-type uridine monomer units for incorporation in 2'-5' RNA: 3'-5' RNA duplexes</b><br>A. D. Gunjal, N. Erande, M. D'Costa and V. A. Kumar                                              | 191–192 |
| <b>A short C-rich PNA fragment capable to form novel G-quadruplex-PNA complexes</b><br>J. Amato, V. Gabelica, N. Borbone, F. Rosu, E. D. Pauw, G. Oliviero, G. Piccialli and L. Mayol                                                    | 167–168 | <b>Structural analysis of Musashi-RNA complex on the basis of long-range structural information</b><br>T. Ohyama, A. Furukawa, T. Mashima, T. Sugiyama, S. Ohgara, T. Yamazaki, T. Imai, H. Okano, T. Nagata and M. Katahira     | 193–194 |
| <b>Human telomere RNA and DNA form an intermolecular G-quadruplex</b><br>Y. Xu, T. Kimura and M. Komiyama                                                                                                                                | 169–170 | <b>Construction of a stable functional ribonucleopeptide complex by the covalent linking method</b><br>M. Fukuda, S. Nakano, K. Tainaka, N. Fujieda and T. Morii                                                                 | 195–196 |
| <b>Orientation of ends of G-quadruplex structure investigated with end-extended oligonucleotides</b><br>Y. Sannohe, K. Sato, A. Matsugami, K. Shinohara, T. Mashimo, M. Katahira and H. Sugiyama                                         | 171–172 | <b>The specific interaction between metal cation and mismatch base pair in duplex RNA</b><br>T. Kozasa, Y. Miyakawa, A. Ono and H. Torigoe                                                                                       | 197–198 |
| <b>Molecular docking study of binding of TMPyP4 to a bimolecular human telomeric G-quadruplex</b><br>Y. Ishikawa and S. Fujii                                                                                                            | 173–174 | <b>Selective recognition of a tetra-amino-acid motif containing phosphorylated tyrosine residue by ribonucleopeptide</b><br>S. Nakano, T. Hasegawa, M. Fukuda, N. Fujieda, K. Tainaka and T. Morii                               | 199–200 |
| <b>Human Telomeric RNA in G-quadruplex Structure</b><br>Y. Xu, K. Kaminaga and M. Komiyama                                                                                                                                               | 175–176 | <b>Development of ribonucleopeptide-based fluorescent sensors for biologically active amines based on the stepwise molding strategy</b><br>K. Tainaka, T. Hasegawa, M. Fukuda, S. Nakano, N. Fujieda and T. Morii                | 201–202 |
| <b>Studies on the influence of inversion of polarity sites on the dG residues glycosidic conformation in quadruplex structures</b><br>V. Esposito, G. Oliviero, A. Pepe, A. Virgilio and A. Galeone                                      | 177–178 | <b>Significance of the N-terminal Histidine-rich Region for the Function of the Human Toll-like Receptor 3 Ectodomain</b><br>T. Tokisue, T. Watanabe, T. Tsujita, S. Nishikawa, T. Hasegawa, T. Seya, M. Matsumoto and K. Fukuda | 203–204 |
| <b>Structural analysis of r(GGA)<sub>4</sub> found in RNA aptamer for bovine prion protein</b><br>A. Matsugami, T. Mashima, F. Nishikawa, K. Murakami, S. Nishikawa, K. Noda, T. Yokoyama and M. Katahira                                | 179–180 | <b>Isolation of RNA aptamers specific for the 3' X tail of HCV</b><br>K. Fukuda, Y. Toyokawa, K. Kikuchi, K. Konno, R. Ishihara, C. Fukazawa, S. Nishikawa and T. Hasegawa                                                       | 205–206 |
| <b>NMR structural study of DNA oligomers containing alkylene crosslinked cyclic 2'-deoxyuridylate dimers</b><br>K. Furuuta, S. Murata, J. Jee, S. Ichikawa, A. Matsuda and C. Kojima                                                     | 181–182 | <b>Subsites for substrate recognition by bacterial ribonuclease P</b><br>A. Fujimoto, S. Suwa, Y. Nagai, Y. Kikuchi and T. Tanaka                                                                                                | 207–208 |
| <b>NMR assignments and the identification of the secondary structure of the anti-retroviral cytidine deaminase</b><br>A. Furukawa, T. Nagata, Y. Habu, R. Sugiyama, F. Hayashi, S. Yokoyama, H. Takaku and M. Katahira                   | 183–184 | <b>Analysis of the interaction between selected RNA-binding peptides and a target RNA containing a bulge and a GNRA-type tetraloop</b><br>S. Horiya, C.-S. Koh, S. Matsufuji and K. Harada                                       | 209–210 |
| <b>Inhibition of human papillomavirus replication by using artificial zinc-finger nucleases</b><br>T. Mino, T. Mori, Y. Aoyama and T. Sera                                                                                               | 185–186 | <b>Fluorescent Ligand as a molecular probe for the RNA structure</b><br>S. Umemoto, J. Zhang, C. Dohno and K. Nakatani                                                                                                           | 211–212 |
| <b>Modulation of endogenous VEGF-A expression under hypoxia by using artificial transcription factors</b><br>T. Mori, J. Sasaki, Y. Aoyama and T. Sera                                                                                   | 187–188 |                                                                                                                                                                                                                                  |         |

|                                                                                                                                                                                                                                    |         |                                                                                                                                                                                                                                              |         |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Identification of RNA Binding Specificity for the TET-family Proteins</b><br>K. Takahama, K. Kino, S. Arai, R. Kurokawa and T. Oyoshi                                                                                           | 213–214 | <b>Synthesis of a perylene diimide derivative having two ferrocene moieties as an electrochemical indicator for human telomeric DNA tetraplex</b><br>R. Fujikawa, S. Watanabe, K. Ohtsuka, S. Sato and S. Takenaka                           | 241–242 |
| <b>Isoenergetic microarray mapping reveals differences in structure between tRNA<sub>i</sub><sup>Met</sup> and tRNA<sub>m</sub><sup>Met</sup> from <i>Lupinus luteus</i></b><br>E. Kierzek, M. Z. Barciszewska and J. Barciszewski | 215–216 | <b>Synthetic Biology for Improved Personalized Medicine</b><br>S. A. Benner, S. Hoshika, M. Sakeda, D. Hutter, N. Leal, Z. Yang and F. Chen                                                                                                  | 243–244 |
| <b>Interactions between antitumor drugs and vault RNA</b><br>T. Mashima, M. Kudo, Y. Takada, A. Matsugami, S. C. B. Gopinath, P. K. R. Kumar and M. Katahira                                                                       | 217–218 | <b>Stability and Mismatch Discrimination of DNA duplexes containing 2,6-Diaminopurine and 2-Thiothymidine Locked Nucleic Acid bases</b><br>C. B. Hughesman, R. F. B. Turner and C. Haynes                                                    | 245–246 |
| <b>Isoenergetic microarray mapping – the advantages of this method in studying the structure of <i>Saccharomyces cerevisiae</i> tRNA<sup>Phe</sup></b><br>M. Jenek and E. Kierzek                                                  | 219–220 | <b>Sensitive DNA probe for photochemical ligation prepared in click chemistry</b><br>T. Ami, H. Matsumura and K. Fujimoto                                                                                                                    | 247–248 |
| <b>Crystal structures of RNA 3'-terminal phosphate cyclase and its complexes with Mg<sup>2+</sup> + ATP, ATP or Mn<sup>2+</sup></b><br>S. Shimizu, M. Ohki, N. Ohkubo, K. Suzuki, M. Tsunoda, T. Sekiguchi and A. Takénaka         | 221–222 | <b>SPR-imaging based assays on an oligonucleotide-array to analyze DNA lesions recognition and excision by repair proteins</b><br>C. Corne, J.-B. Fiche, V. Cunin, A. Buhot, J. Fuchs, R. Calemczuk, A. Favier, T. Livache and D. Gasparutto | 249–250 |
| <b>Lipophilic DNA-conjugates: DNA controlled assembly of liposomes</b><br>U. Jakobsen and S. Vogel                                                                                                                                 | 223–224 | <b>Ion-Exchange Capture of Labeled Pyrrolidinyl Peptide Nucleic Acids for DNA Sequence Determination</b><br>P. Korkaew and T. Vilaivan                                                                                                       | 251–252 |
| <b>DNA Controlled Assembly of Soft Nanoparticles</b><br>U. Jakobsen, A. C. Simonsen and S. Vogel                                                                                                                                   | 225–226 | <b>New substituted aryldiazomethyl labels for high density DNA Chipbased nucleic acid testing</b><br>A. Laurent, M. Kotera, A. Burr, F. Lasnet, T. Martin and A. Laayoun                                                                     | 253–254 |
| <b>Poly (L-lysine)-graft-dextran acts as a nucleic acid chaperone for tetramolecular quadruplex formation</b><br>R. Moriyama, N. Shimada, A. Kano and A. Maruyama                                                                  | 227–228 | <b>Synthesis and Properties of Oligonucleotides Containing Silylated Pyrene Derivatives</b><br>T. Moriguchi, M. Hattori, T. Sekiguchi, M. Ichimura, T. Kato and K. Shinozuka                                                                 | 255–256 |
| <b>New eximer-based tandem systems for SNP detection</b><br>D. S. Novopashina, M. I. Meschaninova, S. A. Kholodar, A. A. Lomzov and A. G. Venyaminova                                                                              | 229–230 | <b>Enhanced amplification of polymerase chain reaction by addition of cosolutes derived from a cellular compatible solute</b><br>K. Koumoto, H. Ochiai and N. Sugimoto                                                                       | 257–258 |
| <b>Design of a fluorescent probe for DNA/RNA imaging</b><br>A. Okamoto, T. Kubota and S. Ikeda                                                                                                                                     | 231–232 | <b>Heat Activatable 3'-modified dNTPs: Synthesis and Application for Hot Start PCR</b><br>I. Koukhareva, H. Haoqiang, J. Yee, J. Shum, N. Paul, R. I. Hogrefe and A. V. Lebedev                                                              | 259–260 |
| <b>Studies of oligodeoxyfluorosides (ODFs) as FRET probes for DNA hybridization</b><br>Y. N. Teo and E. T. Kool                                                                                                                    | 233–234 | <b>Synthesis of novel fluorophores for labelling of oligonucleotides</b><br>A. Singh and R. K. Singh                                                                                                                                         | 261–262 |
| <b>Sequence-selective fluorophore formation by furan-conjugated oligodeoxyribonucleotides</b><br>A. Kobori, J. Morita, M. Ikeda, A. Yamayoshi and A. Murakami                                                                      | 235–236 | <b>Thymidine analogs as potential antiviral agents</b><br>D. Yadav, D. Rai, J. Balzarini, E. De Clercq and R. K. Singh                                                                                                                       | 263–264 |
| <b>Application of partially double-stranded DNA probes to high-throughput SNPs genotyping</b><br>T. Ishii, K. Muraki, N. Shimada, A. Kano, N. Nishida, K. Tokunaga and A. Maruyama                                                 | 237–238 | <b>Design and synthesis of novel oxindoles as potential non-nucleosidic reverse transcriptase inhibitors against HIV</b><br>G. Kumari and R. K. Singh                                                                                        | 265–266 |
| <b>Synthesis of a naphthalene diimide derivative having four ferrocene moieties as an electrochemical DNA hybridization indicator</b><br>S. Sato, M. Tsueda, S. Watanabe, K. Ohtsuka and S. Takenaka                               | 239–240 |                                                                                                                                                                                                                                              |         |

|                                                                                                                                                                                                                         |         |                                                                                                                                                                                                                              |         |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Synthesis and structural analysis of [4.3.0]-bicyclothymidine</b><br>A. Stauffiger and C. J. Leumann                                                                                                                 | 267–268 | <b>Bisphosphonate mRNA cap analog attached to Sepharose for affinity chromatography of decapping enzymes</b><br>S. A. Szczepaniak, J. Jemielity, J. Zuberek, J. Kufel and E. Darzynkiewicz                                   | 295–296 |
| <b>Nucleosides with 1,4-dioxane as sugar moiety</b><br>A. S. Madsen, N. Langkjær and J. Wengel                                                                                                                          | 269–270 | <b>Chemical Models and Their Mechanistic Implications for the Transformation of 6-Cyanouridine 5'-Monophosphate Catalyzed by Orotidine 5'-Monophosphate Decarboxylase</b><br>T.-C. Chien, C.-H. Jen, Y.-J. Wu and C.-C. Liao | 297–298 |
| <b>New conformationally restricted DNA mimics</b><br>T. Højland, B. R. Babu and J. Wengel                                                                                                                               | 271–272 | <b>A versatile synthetic approach for the development of libraries of 5',3'-bis-conjugated oligonucleotides</b><br>G. Di Fabio, C. Coppola, J. D'Onofrio, L. D. Napoli and D. Montesarchio                                   | 299–300 |
| <b>Greener Biocatalytic Approach to the Synthesis of Nucleosides and their Precursors</b><br>S. K. Singh, R. K. Sharma, J. Maity, J. Wengel, V. S. Parmar and A. K. Prasad                                              | 273–274 | <b>Synthesis and properties of oligonucleotides having a chemically stable 2-(trimethylsilyl)benzoyl group</b><br>K. Yamada, H. Taguchi, A. Ohkubo, K. Seio and M. Sekine                                                    | 301–302 |
| <b>Synthesis of double-headed nucleosides with the additional nucleobase in the 5'-C-position</b><br>C. Andersen, P. K. Sharma, M. S. Christensen, N. S. Pedersen and P. Nielsen                                        | 275–276 | <b>Synthesis of a new internucleosidic linkage: the Boronucleotides</b><br>D. Luvino, C. Baraguey, M. Smietana and J.-J. Vasseur                                                                                             | 303–304 |
| <b>Efficient Preparation of xanthine-containing oligodeoxynucleotide from oxanine-containing oligodeoxynucleotide, catalyzed by N<sup>ε</sup>-acetyl-L-histidine</b><br>A. Doi, S. P. Pack, T. Kodaki and K. Makino     | 277–278 | <b>Design and Chemical Synthesis of a 2',4'-BNA Probe for 7,8-Dihydro-8-oxoguanine-containing Double Stranded DNA</b><br>T. Miyoshi, T. Kodama, S. Obika and T. Imanishi                                                     | 305–306 |
| <b>Development of a novel selective detection method for 5-formyl-2'-deoxyuridine by fluorescent derivatization</b><br>W. Hirose, K. Sato and A. Matsuda                                                                | 279–280 | <b>Synthesis of C8-N-Acetylarylamine 2'-dG-adducts and their sitespecifically incorporation into oligonucleotides</b><br>S. Krüger and C. Meier                                                                              | 307–308 |
| <b>Synthesis of 3'-deoxyapionucleoside triphosphates and their incorporation into DNA by DNA polymerase</b><br>M. Kataoka, K. Sato and A. Matsuda                                                                       | 281–282 | <b>Synthesis and properties of covalently linked antiparallel and parallel DNA duplexes</b><br>T. Ono, Y. Saotome, I. Okamoto and A. Ono                                                                                     | 309–310 |
| <b>Use of DNA and Click chemistries to synthesize combinatorial libraries of galactosyl-phosphodiester clusters</b><br>F. Morvan, G. Pourceau, A. Meyer and J.-J. Vasseur                                               | 283–284 | <b>Some Observations on Detritylation in Solid-Phase Oligonucleotide Synthesis</b><br>S. Millar, H. (Tony) Yan, Y. S. Sanghvi and R. Pon                                                                                     | 311–312 |
| <b>Synthesis of nucleoside phosphorothio-, phosphorodithio- and phosphoroselenoate diesters via oxidative esterification of the corresponding H-phosphonate analogues</b><br>M. Kalek, A. Bartoszewicz and J. Stawinski | 285–286 | <b>Foldamers derived from nucleoside beta-amino acids: a new twist on the DNA helix</b><br>R. Cosstick, R. Threlfall and J. Fisher                                                                                           | 313–314 |
| <b>Synthesis and biochemical studies of tetraphosphate 5' mRNA cap analogs bearing bisphosphonate modification</b><br>A. Rydzik, J. Zuberek, J. Kowalska, M. Lukaszewicz, E. Darzynkiewicz and J. Jemielity             | 287–288 | <b>2'-Methylseleno modified oligoribonucleotides synthesized for X-ray crystallography</b><br>H. Moroder, B. Puffer and R. Micura                                                                                            | 315–316 |
| <b>The first examples of mRNA cap analogs bearing boranophosphate modification</b><br>J. Kowalska, J. Zuberek, Z. M. Darzynkiewicz, M. Lukaszewicz, E. Darzynkiewicz and J. Jemielity                                   | 289–290 | <b>Chiral Phosphonate Internucleotide Linkage: A Promising Modification for Chimeric Oligonucleotides?</b><br>R. Liboska, M. Petrová, R. Pohl, M. Buděšínský, V. Hurychová and I. Rosenberg                                  | 317–318 |
| <b>m<sup>7</sup>GTP<math>\alpha</math>S is a strong and stable inhibitor of cap-dependent translation</b><br>J. Kowalska, M. Lukaszewicz, J. Zuberek, M. Ziemniak, M. Strenkowska, E. Darzynkiewicz and J. Jemielity    | 291–292 | <b>Incorporation of 3'-S-phosphorothiolates into RNA: potential applications in RNAi</b><br>J. Gaynor, J. Fisher, B. Campbell and R. Cosstick                                                                                | 319–320 |
| <b>Synthesis of a Photoresponsive <math>\alpha</math>-Dideoxyuridine Triphosphate Derivative</b><br>Y. Nitta, M. Ogino, Y. Yoshimura and K. Fujimoto                                                                    | 293–294 |                                                                                                                                                                                                                              |         |

|                                                                                                                                                                                                                                                                                                       |         |                                                                                                                                                                                                                                                             |         |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Optimized Fluoride Ions Treatment for Release of Base-Sensitive 2'-O-modified Oligoribonucleotides from Various Solid Supports</b><br>T. Lavergne, F. Debart and J.-J. Vasseur                                                                                                                     | 321–322 | <b>DNA Glue: 1-, 2- and 4-Ethynylpyrenes in the Structure of Twisted Intercalating Nucleic Acids (TINAs), DNA Duplexes/Triplexes and Interstrand Excimer Formation</b><br>V. V. Filichev, I. V. Astakhova, A. D. Malakhov, V. A. Korshun and E. B. Pedersen | 347–348 |
| <b>Chemical synthesis of RNA including 5-taurinomethyluridine and 5-taurinomethyl-2-thiouridine</b><br>T. Ogata and T. Wada                                                                                                                                                                           | 323–324 | <b>BODIPY-modified uridines as potential fluorescent probes for nucleic acids that are recognized by DNA-polymerases</b><br>T. Ehrenschwender, C. Wanninger-Weiß and H.-A. Wagenknecht                                                                      | 349–350 |
| <b>Development of new methods for the synthesis of RNA oligonucleotides having baselabile functional groups</b><br>A. Ohkubo, Y. Noma, Y. Kuwayama, H. Tsunoda, K. Seio and M. Sekine                                                                                                                 | 325–326 | <b>Cholesterol-Linked Pyrene Excimer Molecular Beacon with Enhanced Cell Permeability</b><br>H. S. Jeong, Y. J. Seo, E.-K. Bang, G. T. Hwang, J. H. Jung, S. K. Jang and B. H. Kim                                                                          | 351–352 |
| <b>Synthesis and properties of 2'-modified-4'-thioRNA</b><br>M. Takahashi, N. Minakawa and A. Matsuda                                                                                                                                                                                                 | 327–328 | <b>Fluorogenic probe triggered by reduction for nucleic acids sensing</b><br>K. Furukawa, H. Abe, J. Wang, K. Oki, M. Uda, S. Tsuneda and Y. Ito                                                                                                            | 353–354 |
| <b>Synthesis and properties of oligonucleotides containing 4'-selenoribonucleosides</b><br>Y. Inagaki, N. Minakawa and A. Matsuda                                                                                                                                                                     | 329–330 | <b>Intracellular mRNA imaging with a hybridization sensitive fluorescent nucleotide</b><br>T. Kubota, S. Ikeda and A. Okamoto                                                                                                                               | 355–356 |
| <b>A convenient approach towards 2'-O-modified RNA-oligonucleotides on solid support using universal nucleosides</b><br>J. Haas and J. W. Engels                                                                                                                                                      | 331–332 | <b>Fluorometric sensing of conformational switching of DNA; The use of fluorescence labeled C8-alkylamino substituted 2'-deoxyguanosine</b><br>Y. Saito, K. Matsumoto, S. S. Bag, A. Misawa and I. Saito                                                    | 357–358 |
| <b>Stereocontrolled synthesis of backbone-modified oligonucleotides via diastereopure H-phosphonate intermediates</b><br>N. Iwamoto, N. Oka and T. Wada                                                                                                                                               | 333–334 | <b>Design of an efficient self-quenched molecular beacon for SNPs genotyping</b><br>Y. Saito, K. Numajiri, S. S. Bag, E. Mizuno and I. Saito                                                                                                                | 359–360 |
| <b>Stereocontrolled synthesis of oligonucleoside phosphorothioates and PO/PS-chimeric oligonucleotides by using oxazaphospholidine derivatives</b><br>N. Oka, M. Yamamoto, T. Sato and T. Wada                                                                                                        | 335–336 | <b>Design of an ultimate quencher free molecular beacon containing pyrrolocytidine-guanine base pair</b><br>Y. Saito, Y. Shinohara, S. S. Bag, Y. Takeuchi, K. Matsumoto and I. Saito                                                                       | 361–362 |
| <b>Towards Fluorous-Phase Synthesis of Nucleoside <math>\beta</math>-Peptides</b><br>R. Threlfall, R. Cosstick and T. Wada                                                                                                                                                                            | 337–338 | <b>Targeting specific gene by alkylating pyrrole-imidazole polyamides</b><br>M. Minoshima, J. Chou, S. Lefebvre, T. Bando, K. Shinohara, J. M. Gottesfeld and H. Sugiyama                                                                                   | 363–364 |
| <b>Translesion synthesis across the (6-4) photoproduct and its Dewar valence isomer by the Y-family and engineered DNA polymerases</b><br>J. Yamamoto, D. Loakes, C. Masutani, S. Simmyo, K. Urabe, F. Hanaoka, P. Holliger and S. Iwai                                                               | 339–340 | <b>Sequence-specific alkylation of DNA by pyrrole-imidazole polyamides through cooperative interaction</b><br>G. Kashiwazaki, T. Bando and H. Sugiyama                                                                                                      | 365–366 |
| <b>Thiophosphoramidites scale up and their use in the synthesis of phosphorodithioate linkages</b><br>X. Yang                                                                                                                                                                                         | 341     | <b>Site-specific modification by functionality-transfer oligonucleotide with the photo-inducible reactivity</b><br>K. Onizuka, Y. Taniguchi, T. Nishioka and S. Sasaki                                                                                      | 367–368 |
| <b>Novel Advanced Universal Solid Supports for Oligonucleotide Synthesis</b><br>A. P. Guzaev                                                                                                                                                                                                          | 343     | <b>Photochromic Nucleobase: Reversible Photoisomerization, Photochemical Properties and Photoregulation of hybridization</b><br>S. Ogasawara and M. Maeda                                                                                                   | 369–370 |
| <b>Synthesis of four colors fluorescently labelled 3'-O-blocked nucleotides with fluoride cleavable blocking group and linker for array based Sequencing-by-Synthesis applications</b><br>D. C. Knapp, A. Keller, J. D'Onofrio, A. Lubys, S. Serva, A. Kurg, M. Remm, M. Kwiatkowski and J. W. Engels | 345–346 |                                                                                                                                                                                                                                                             |         |

|                                                                                                                                                                                                        |         |                                                                                                                                                                                                                  |         |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| Synthesis and incorporation of fluorescent C-nucleoside analogue into oligodeoxyribonucleotides<br>H. Ozaki, T. Kawai, M. Kuwahara and H. Sawai                                                        | 371–372 | RNA containing pyrrolocytidine base analogs: good binding affinity and fluorescence that responds to hybridization<br>A. S. Wahba, M. J. Damha and R. H. E. Hudson                                               | 399–400 |
| Enzymatic Synthesis of multi Spin-labeled DNA<br>S. Obeid, M. Yulikov, G. Jeschke and A. Marx                                                                                                          | 373–374 | Exceptional Fluorescence and Hybridization Properties of a Phenylpyrrolocytosine in Peptide Nucleic Acid<br>F. Wojciechowski and R. H. E. Hudson                                                                 | 401–402 |
| Synthesis and Relaxivities of Oligonucleotides with TEMPO-labeled Sugar Moiety<br>M. Okazaki, Y. Sato, S. Karasawa and N. Koga                                                                         | 375–376 | Synthesis of RNA Dimer and Trimer Blocks and Their Uses<br>S. W. Kim, A. Kayushin, M. Kim, A. Miroshnikov and K. E. Jung                                                                                         | 403     |
| Structure–activity relationship of an antisense oligonucleotide–two Cu(II) complex conjugate as an artificial ribonuclease<br>D. Mukoguchi, S. Sakamoto, H. Takayama, M. Kitamura and H. Inoue         | 377–378 | Easy method for the synthesis of labeled oligonucleotides<br>D. Heindl, D. Kessler, A. Schube, W. Thuer and A. Giraut                                                                                            | 405–406 |
| Hydrolysis of Single Linkage in Long DNA using Oligonucleotide-Multiphosphonate Conjugates and Ce <sup>IV</sup> /EDTA<br>Y. Suzuki, T. Lönnberg and M. Komiyama                                        | 379–380 | Synthesis of oligonucleotides terminally modified by bulky and negatively charged substituents and their hybridization with RNA targets<br>K. Seio, Y. Takaku, K. Miyazaki, S. Kurohagi, A. Ohkubo and M. Sekine | 407–408 |
| New Methodology to Search for DNA Binding Ligands Based on Duplex DNA-Templated Click Chemistry<br>S. Imoto, T. Hirohama and F. Nagatsugi                                                              | 381–382 | Folding pathways of hybrid-1 and hybrid-2 G-quadruplex structures<br>T. Mashimo, Y. Sannohe, H. Yagi and H. Sugiyama                                                                                             | 409–410 |
| DNA Conjugation by Staudinger Ligation<br>S. H. Weisbrod, A. Baccaro and A. Marx                                                                                                                       | 383–384 | Mechanism of DNA elongation during <i>de novo</i> DNA synthesis<br>X. Liang, T. Kato and H. Asanuma                                                                                                              | 411–412 |
| A new linker group for functionalization of nucleic acids<br>V. Svetlana, K. Dmitrii, B. Boris, A. Tatiana, K. Nina and S. Vladimir                                                                    | 385–386 | Thermodynamics of DNA structures under molecular crowding conditions with neutral and positive charged cosolutes<br>D. Miyoshi, K. Nakamura, S. Muhuli, H. T. Karimata and N. Sugimoto                           | 413–414 |
| Synthesis of oligonucleotide conjugated pyrrolepolyamide- and pyrrole-imidazole polyamide-2'-deoxyguanosine hybrids as novel gene expression control compounds<br>E. Kawashima, Y. Ohba and K. Kamaike | 387–388 | Mechanism of chiral-selective tRNA aminoacylation and the origin of amino acid homochirality<br>K. Tamura                                                                                                        | 415–416 |
| Circular DNA Formation on the Bimolecular Triplex of Anthracene-modified Oligonucleotide Conjugate<br>P. Arslan, T. Ihara and A. Jyo                                                                   | 389–390 | Effect of Locked Nucleic Acid (LNA) modification on Hybridization Kinetics of DNA duplex<br>A. Arora, H. Kaur, J. Wengel and S. Maiti                                                                            | 417–418 |
| Synthesis of the Peptide Nucleic Acid (PNA) Incorporating 2-Amino-6-vinylpurine Derivative and Evaluation of the Reactivity<br>S. Imoto, A. Shimotazawa, T. Hirohama, M. Goto and F. Nagatsugi         | 391–392 | Promotion of Triplex Formation by 2'-O,4'-C-Aminomethylene Bridged Nucleic Acid (2',4'-BNA <sup>NC</sup> ) Modification<br>K. Sasaki, S. M. A. Rahman, S. Obika, T. Imanishi and H. Torigoe                      | 419–420 |
| Construction of an aminoxy derivative for RNA and DNA labeling<br>Y. Komatsu, N. Kojima, T. Takebayashi, A. Mikami, M. Sugino and E. Ohtsuka                                                           | 393–394 | Photoinduced excess electron injection into DNA duplexes containing mismatched base pairs<br>T. Ito, A. Kondo, A. Hayashi, T. Uchida, K. Tanabe and S. Nishimoto                                                 | 421–422 |
| Effective Synthesis of Photosensitive Oligodeoxynucleotides<br>M. Ogino, Y. Taya and K. Fujimoto                                                                                                       | 395–396 | Development of template-directed reversible DNA photocrosslinking<br>Y. Yoshimura, G. Ozaki and K. Fujimoto                                                                                                      | 423–424 |
| RNA containing pyrrolocytidine base analogs: increased binding affinity and fluorescence that responds to hybridization<br>A. S. Wahba, M. J. Damha and R. H. E. Hudson                                | 397–398 |                                                                                                                                                                                                                  |         |

|                                                                                                                                                                                                                                 |         |                                                                                                                                                                                                                                          |         |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Thermodynamic, Counterion and Hydration Effects for the Incorporation of Locked Nucleic Acid (LNA) Nucleotides in Duplex</b><br>H. Kaur, A. Arora, J. Wengel and S. Maiti                                                    | 425–426 | <b>X-Ray Crystallographic Identification of Bisulfite-Uracil Adduct as Sodium 5,6-Dihydrouracil 6-Sulfonate</b><br>M. Kaneda, Y. Hashimoto and H. Hayatsu                                                                                | 451–452 |
| <b>On the Stereochemistry of C3'-O-P-CH<sub>2</sub>-O-C4'' Phosphonate Internucleotide Bond, a Phosphate Isostere</b><br>Z. Točík, M. Buděšínský, I. Barvík and I. Rosenberg                                                    | 427–428 | <b>Effect of backbone-modification of oligodeoxyribonucleic acid on primer extension reactions</b><br>M. Kuwahara, S. Minezaki, J. Nagashima, H. Ozaki and H. Sawai                                                                      | 453–454 |
| <b>Temperature-dependent changes of RNA secondary structures monitored by <sup>19</sup>F-NMR spectroscopy</b><br>D. Graber and R. Micura                                                                                        | 429     | <b>Towards the Replication of xDNA, a Size-expanded Unnatural Genetic System</b><br>A. T. Krueger, H. Lu, T. Højland, H. Liu, J. Gao and E. T. Kool                                                                                      | 455–456 |
| <b>Effect of Linker Length on DNA Duplexes Containing a Mismatched O<sup>6</sup>-2'-Deoxyguanosine-Alkyl Interstrand Cross-Link</b><br>J. D. Booth, S. P. Murphy, A. M. Noronha and C. J. Wilds                                 | 431–432 | <b>Sequences around the unnatural base pair in DNA templates for efficient replication</b><br>M. Kimoto, R. Kawai, T. Mitsu, S. Yokoyama and I. Hirao                                                                                    | 457–458 |
| <b>Reactivity of Thymine Doublet in Single Strand DNA with Osmium Reagent</b><br>A. Nomura and A. Okamoto                                                                                                                       | 433–434 | <b>Radiolytic Ligation of Oligodeoxynucleotides Possessing Disulfide Bond</b><br>K. Tanabe, E. Kuraseko, E. Matsumoto, H. Yamada, T. Ito and S. Nishimoto                                                                                | 459–460 |
| <b>The reaction of cytosine with bisulfite by base flipping from the duplex</b><br>Y. Oka, T. Peng, F. Takei and K. Nakatani                                                                                                    | 435–436 | <b>Regulation of Protein-protein Interaction <i>via</i> Assembly of Coiled-coil Domain</b><br>K. Suzuki, T. Mizuno and T. Tanaka                                                                                                         | 461     |
| <b>Stereoselective formation of a cyclobutane pyrimidine dimer by using N<sup>4</sup>-acetyl protection of the cytosine base</b><br>K. Nishiguchi, J. Yamamoto and S. Iwai                                                      | 437–438 | <b>Comparison of the chemical properties of a novel amino-linker with various amino modifications</b><br>N. Kojima, T. Takebayashi, A. Mikami, M. Sugino, E. Ohtsuka and Y. Komatsu                                                      | 463–464 |
| <b>Oligonucleotides damaged with Carcinogenic Aromatic Amines at the C8- and N<sup>6</sup>-Position of 2'-Deoxyadenosine</b><br>M. I. Jacobsen and C. Meier                                                                     | 439–440 | <b>DNA Strand Exchange on Liposome Surfaces</b><br>K. Frykholm, F. B. Bombelli, B. Nordén and F. Westerlund                                                                                                                              | 465     |
| <b>Reactivity of oxanine: efficient fabrication of DNA microarray by using oxanine-containing DNA oligomer as probe molecule</b><br>S. P. Pack, N. K. Kamisetty, M. Nonogawa, T. Kodaki and K. Makino                           | 441–442 | <b>Direct Preparation of Sticky-Ended Duplexes within PCR by Using Caged Primers</b><br>K. Tanaka, H. Katada, N. Shigi, A. Kuzuya and M. Komiyama                                                                                        | 467–468 |
| <b>Analysis for complexity of clustered DNA damage generated by heavy ion beams</b><br>H. Terato, H. Watari, Y. Shimazaki, R. Hirayama, Y. Furusawa and H. Ide                                                                  | 443–444 | <b>Efficient PCR amplification by an unnatural base pair system</b><br>M. Kimoto, R. Kawai, T. Mitsui, S. Yokoyama and I. Hirao                                                                                                          | 469–470 |
| <b>Base recognition abilities of 2'-deoxynucleoside N-oxide derivatives and their substrate specificity in enzyme reaction</b><br>H. Tsunoda, T. Kudo, A. Ohkubo, K. Seio and M. Sekine                                         | 445–446 | <b>Azide-alkyne “click” reaction performed on oligonucleotides with the universal nucleoside 7-octadiynyl-7-deaza-2'-deoxyinosine</b><br>X. Ming, P. Leonard, D. Heindl and F. Seela                                                     | 471–472 |
| <b>pH effect on the one-electron photooxidation of 5-methylcytosine with naphthoquinone sensitizer</b><br>H. Yamada, K. Tanabe, T. Ito and S. Nishimoto                                                                         | 447–448 | <b>Temperature-gradient dependant detection of target DNA oligomers using DNA-immobilized open tubular capillary column</b><br>N. K. Kamisetty, S. P. Pack, K. C. Devarayapalli, M. Nonogawa, T. Kodaki and K. Makino                    | 473–474 |
| <b>Synthesis, Biophysical and Repair Studies of O<sup>6</sup>-2'-Deoxyguanosine Adducts by <i>Escherichia coli</i> OGT</b><br>N. M. Schoonhoven, S. P. Murphy, D. K. O'Flaherty, A. M. Noronha, M. J. Kornblatt and C. J. Wilds | 449–450 | <b>Toward the discovery of new antifungal agents: The design and validation of a novel 2'-P-RNA probe and high throughput screening assay against 2'-phosphotransferase Tpt1p</b><br>J. G. Lackey, D. Ron, M. J. Damha and H. P. Harding | 475–476 |

|                                                                                                                                                                                        |         |                                                                                                                                                                                                                                                  |         |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>New Strategies for DNA Polymerase Library Screening</b><br>R. Kranaster and A. Marx                                                                                                 | 477–478 | <b>Nanocircular RNAs for RNA interference</b><br>H. Abe, N. Abe, M. Harada, S. Tsuneda and Y. Ito                                                                                                                                                | 505–506 |
| <b>Codon based mutagenesis using trimer phosphoramidites</b><br>J. Randolph, A. Yagodkin, M. Lamaitre, A. Azhayev and H. Mackie                                                        | 479     | <b>MicroRNAs as biomarkers in Tomato Leaf Curl Virus (ToLCV) disease</b><br>A. R. Naqvi, N. R. Choudhury, Q. M. Rizwanul Haq and S. K. Mukherjee                                                                                                 | 507–508 |
| <b>Intracellular small RNA-agarose: Preparation and application for the analysis of proteins interacted with small RNA</b><br>M. Fujino and S. Izuta                                   | 481–482 | <b>Micro RNA like inhibition of HIV-1 replication</b><br>K. Kato, Y. Habu and H. Takaku                                                                                                                                                          | 509–510 |
| <b>Site-specific gene manipulation of fluorescent proteins using artificial restriction DNA cutter</b><br>H. Katada, Y. Ren, N. Shigi and M. Komiyama                                  | 483–484 | <b>The HIV-2 TAR RNA domain as a potential source of viral-encoded miRNA. A reconnaissance study</b><br>K. J. Purzycka and R. W. Adamiak                                                                                                         | 511–512 |
| <b>Oligonucleotides are potent antioxidants acting mainly as metal-ion chelators</b><br>B. Fischer and E. Zobel                                                                        | 485–486 | <b><i>In vitro</i> selection of RNA aptamer to hemin</b><br>M. Liu, T. Kagahara, H. Abe and Y. Ito                                                                                                                                               | 513–514 |
| <b>RNA aptamers specifically interact with the Fc region of mouse immunoglobulin G</b><br>N. Sakai, H. Masuda, J. Akitomi, H. Yagi, Y. Yoshida, K. Horii, M. Furuichi and I. Waga      | 487–488 | <b>Photodynamic antisense regulation of mRNA having a point mutation with psoralen-conjugated oligonucleotide</b><br>M. Higuchi, A. Yamayoshi, A. Kobori and A. Murakami                                                                         | 515–516 |
| <b>A multisubstrate deoxyribonucleoside kinase from plants</b><br>A. R. Clausen, L. Girandon, W. Knecht, S. Survery, E. Andreasson, B. Munch-Petersen and J. Piškur                    | 489–490 | <b>Self-catalyzing functions of DNA</b><br>H. Maeda, M. Ikeguchi, T. Niitu and N. Minoura                                                                                                                                                        | 517–518 |
| <b>Influence of 3'-azido-2',3'-dideoxyguanosine treatment on amounts of TERT and POT1 in human HL60 cells</b><br>X. Liu, T. Ogawara, Y. Kurashina, M. Saneyoshi and T. Yamaguchi       | 491–492 | <b>Molecular crowding effect on metal ion binding properties of the hammerhead ribozyme</b><br>S.-i. Nakano, Y. Kitagawa, H. T. Karimata and N. Sugimoto                                                                                         | 519–520 |
| <b>Isolation and characterization of RNA aptamers specific for the HCV minus-IRES domain I</b><br>K. Konno, S. Fujita, M. Iizuka, S. Nishikawa, T. Hasegawa and K. Fukuda              | 493–494 | <b>DNAzymes and ribozymes carrying 2'-C-methyl nucleotides</b><br>R. Pontiggia, L. Robaldo, J. Montserrat and A. Iribarren                                                                                                                       | 521–522 |
| <b>Expanding substrate specificity of nucleoside 2'-deoxyribosyltransferase</b><br>P. A. Kaminski, L. Dugué and S. Pochet                                                              | 495–496 | <b>Mutation analysis of the base-pair connecting two functional modules in the DSL ribozyme</b><br>J. Ishikawa, H. Furuta and Y. Ikawa                                                                                                           | 523–524 |
| <b>Purification of eukaryotic translation factors from wheat germ for reconstitution of protein synthesis</b><br>H. Nagano, S. Sugihara, H. Takagi, T. Ogasawara, Y. Endo and K. Takai | 497–498 | <b>Effective cleavage of structured RNAs by tandems of 10-23 DNAzymes with 3'-modified oligo(2'-O-methylribonucleotide)-effectors</b><br>A. Fokina, D. Novopashina, M. Meschaninova, M. Vorobjeva, M. Zenkova, J.-C. Francois and A. Venyaminova | 525–526 |
| <b>Suppression of bcr-abl mRNA by Chemically Modified siRNA</b><br>S. Murao, I. Diala and M. Fujii                                                                                     | 499–500 | <b>Simple and Rapid Colorimetric Detection of Low-Weight Molecules Using Aptazymes in Combination with Noncrosslinking Gold Nanoparticle Aggregation</b><br>A. Ogawa and M. Maeda                                                                | 527–528 |
| <b>Tc-DNA modified siRNA</b><br>D. Ittig, D. Schümperli and C. J. Leumann                                                                                                              | 501–502 | <b>Design and synthesis of peptide-oligonucleotide conjugates as potential artificial ribonucleases</b><br>I. Serpokyrova, L. Koroleva, N. Svischeva, D. Novopashina and V. Silnikov                                                             | 529–530 |
| <b>Synthesis and silencing properties of siRNAs possessing lipophilic groups at their 3'-termini</b><br>Y. Ueno, K. Kawada, A. Shibata, K. Yoshikawa, Y. Wataya and Y. Kitade          | 503–504 |                                                                                                                                                                                                                                                  |         |

|                                                                                                                                                                                                                                                                         |         |                                                                                                                                                                                                                                |         |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Kinetics of dCTP incorporation opposite to 7,8-dihydro-8-oxoguanine with different 5' nearest neighbors by yeast polymerase <math>\eta</math></b><br>C. W. Yung, D. Loakes, S. Arimoto, K. Negishi and T. Negishi                                                    | 531–532 | <b>Synthetic study of muraymycins using Ugi-four component reaction</b><br>T. Tanino, S. Hirano, S. Ichikawa and A. Matsuda                                                                                                    | 557–558 |
| <b>6-Aryl- and 6-Heteroarylpurines via Cyclotrimerization</b><br>P. Turek, M. Hocek and M. Kotora                                                                                                                                                                       | 533–534 | <b>Total syntheses of a North methanocarba Puromycin analog and its dinucleotide derivative</b><br>B. Y. Michel and P. Strazewski                                                                                              | 559–560 |
| <b>Synthesis and some transformation of acyclic nucleotide phosphonate analogues with triple bond</b><br>V. K. Brel                                                                                                                                                     | 535–536 | <b>Regioselective glycosylation of 7-azapteridines and conversion of the 7-azapteridine nucleosides into 6-azapurine nucleosides</b><br>T. Nagamatsu, J. Ma and S. Akiba                                                       | 561–562 |
| <b>Prolinol-Based Nucleoside Phosphonic Acids: Synthesis and Properties</b><br>V. Vaněk, M. Buděšínský, R. Liboska, V. Hurychová and I. Rosenberg                                                                                                                       | 537–538 | <b>Improved Synthesis of Nucleoside Diphosphate Glycopyranoses</b><br>T. Zismann and C. Meier                                                                                                                                  | 563–564 |
| <b>A step further in the SATE mononucleotide prodrug approach</b><br>S. Peyrottes, A.-L. Villard, G. Coussot, P. Augustijns, I. Lefebvre, A.-M. Aubertin, G. Gosselin and C. Périgaud                                                                                   | 539–540 | <b>Exploring synthetic routes to nucleoside alkynylphosphonates</b><br>M. Meurillon, S. Peyrottes and C. Périgaud                                                                                                              | 565–566 |
| <b>Coupled biocatalysts applied to the synthesis of nucleosides</b><br>R. Medici, M. T. Porro, E. Lewkowicz, J. Montserrat and A. M. Iribarren                                                                                                                          | 541–542 | <b>Reactive oxygen species generation through NADH oxidation by pterin derivatives</b><br>M. Nonogawa, T. Arai, N. Endo, S. P. Pack, T. Kodaki and K. Makino                                                                   | 567–568 |
| <b>What are the consequences of freezing the anomeric effect in nucleosides?</b><br>V. E. Marquez, G. Sun, M. A. Siddiqui, Y.-C. Lee, J. J. Barchi, I. V. Filippov, N. A. Landsman and J. A. Kelley                                                                     | 543–544 | <b>Synthesis of Nucleoside 5'-S-methylphosphonates and Related Compounds</b><br>I. Kóšiová and I. Rosenberg                                                                                                                    | 569–570 |
| <b>First enantioselective synthesis of (–)-neplanocin B</b><br>N. Hamon, J.-P. Uttaro, C. Mathé and C. Périgaud                                                                                                                                                         | 545–546 | <b>“Reverse Fleximers”: Introduction of a series of 5-substituted carbocyclic uridine analogues</b><br>J. M. Sadler, O. Ojewoye and K. L. Seley-Radtke                                                                         | 571–572 |
| <b>Synthesis of some derivatives of 3'-C-methyl pyrimidine nucleosides</b><br>M. Aljarah, C. Mathé and C. Périgaud                                                                                                                                                      | 547–548 | <b>Synthesis of <i>N</i>-1-alkyl analogues of cyclic inosine diphosphate ribose (cIDPR) by a new solid phase approach</b><br>G. Oliviero, S. D'Errico, N. Borbone, J. Amato, V. Piccialli, M. Varra, G. Piccialli and L. Mayol | 573–574 |
| <b>A one step synthetic approach to L-pyrimidine nucleosides using Natural Phosphate coated with potassium iodide as catalyst</b><br>H. B. Lazrek, L. Baddi, D. Ouzebila, M. Smietana and J.-J. Vasseur                                                                 | 549–550 | <b>Total synthesis of a <i>xylo</i>-Puromycin analog</b><br>B. Y. Michel, K. S. Krishnakumar and P. Strazewski                                                                                                                 | 575–576 |
| <b>Synthesis of functionalized <math>\delta</math>-amino alcohol stereoisomers with a cyclopentane skeleton. Hydroxylated azidocarbanucleoside precursors</b><br>L. Kiss, E. Forró, R. Sillanpää and F. Fülöp                                                           | 551–552 | <b>Pyrrolidine analogues of nucleosides and nucleotides</b><br>D. Rejman, R. Pohl, S. Kovačková, P. Kočalka, A. Švenková, H. Šanderová, L. Krásný and I. Rosenberg                                                             | 577–578 |
| <b>Design, Synthesis And Evaluation Of Constrained Methoxyethyl (cMOE) and Constrained Ethyl (cEt) Nucleoside Analogs</b><br>P. P. Seth, A. Siwkowski, C. R. Allerson, G. Vasquez, S. Lee, T. P. Prakash, G. Kinberger, M. T. Migawa, H. Gaus, B. Bhat and E. E. Swayze | 553–554 | <b>New Efficient Synthesis of Nucleoside Diphosphate Glycopyranoses</b><br>S. Wolf and C. Meier                                                                                                                                | 579–580 |
| <b>Synthesis and Conformational Analysis of Novel 2', 3'-Didehydro-2', 3'-dideoxy-4'-selenonucleosides</b><br>Y. J. Ko, W. J. Choi, J. H. Jun, L. X. Zhao and L. S. Jeong                                                                                               | 555–556 | <b>Highly <math>\beta</math>-selective, N-iodosuccinimide-mediated nucleosidation to bicyclo- and tricyclo-nucleosides</b><br>S. Luisier, P. Silhar and C. J. Leumann                                                          | 581–582 |
|                                                                                                                                                                                                                                                                         |         | <b>New and efficient Synthesis of Nucleoside Polyphosphates and Nucleoside Monophosphate Sugars</b><br>S. Warnecke and C. Meier                                                                                                | 583–584 |
|                                                                                                                                                                                                                                                                         |         | <b>Synthesis of 5-ethynyl-1-<math>\beta</math>-D-ribofuranosyl-1<i>H</i>-[1,2,3]triazole-4-carboxylic acid amide (isosteric to EICAR) and its derivatives</b><br>T. Ostrowski and J. Zeidler                                   | 585–586 |

|                                                                                                                                                                                                                                                      |         |                                                                                                                                                                                                                                                                                                                                |         |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Piperidine nucleosides and nucleotides</b><br>S. Kovačková, O. Pačes, M. Dračinský, I. Rosenberg and D. Rejman                                                                                                                                    | 587     | <b>Synthesis and antiviral evaluation of a seven-membered sugar ring nucleoside analog, 9-(5-deoxy-β-D-<i>allo</i>-septanosyl)-adenine</b><br>G. Sizun, J. Griffon, L. Griffe, D. Dukhan, R. Storer, J.-P. Sommadossi, A. G. Loi, C. Musiu, B. Poddesu, A. Cadeddu, M. Fanti, N. Boscu, F. Bassetti, M. Liuzzi and G. Gosselin | 611–612 |
| <b>Novel nucleotide phosphonate analogues with 1,2-oxaphosphol-3-ene ring skeleton</b><br>V. K. Brel                                                                                                                                                 | 589–590 |                                                                                                                                                                                                                                                                                                                                |         |
| <b>From Ribonucleoside 5'-Aldehydes to Ribonucleoside 5'-C-phosphonates as Building Blocks for Oligonucleotide Synthesis</b><br>M. Petrová, Š. Králíková, M. Buděšínský and I. Rosenberg                                                             | 591–592 | <b>Synthesis of Novel Cyclopropyl Nucleoside Analogues as Potential Antiherpetic Agent</b><br>J.-A. Kang, A.-Y. Park, H.-R. Lee, W. H. Kim, P. Chun, L. S. Jeong and H. R. Moon                                                                                                                                                | 613–614 |
| <b>Design and Synthesis of 1-(β-D-Ribofuranosyl)imidazo[4,5-c]pyrazoles as 5:5 Bicyclic Analogs of Purine Nucleosides</b><br>T.-C. Chien, J. C. Drach and L. B. Townsend                                                                             | 593–594 | <b>Carbocyclic L-Nucleoside Analogs as Potential Antiviral Agents</b><br>S. Jessel and C. Meier                                                                                                                                                                                                                                | 615–616 |
| <b>Synthesis and antiviral evaluation of 7-fluoro-7-deaza-2-aminopurine nucleoside derivatives</b><br>F. Leroy, D. Chaves, D. Dukhan, R. Storer, J.-P. Sommadossi, A. G. Loi, A. Cadeddu, M. Fanti, N. Boscu, F. Bassetti, M. Liuzzi and G. Gosselin | 595–596 | <b>Synthesis and antiviral evaluation of 4-fluoropyrazole-3-carboxamide nucleoside derivatives</b><br>F. Leroy, D. Chaves, D. Dukhan, R. Storer, J.-P. Sommadossi, A. Giulia Loi, A. Cadeddu, M. Fanti, N. Boscu, F. Bassetti, M. Liuzzi and G. Gosselin                                                                       | 617–618 |
| <b>New depot forms of AZT and 3TC based on their phosphonate derivatives: anti-HIV activity and pharmacokinetic parameters</b><br>M. K. Kukhanova, M. V. Jasco, A. V. Shipitzyn, D. V. Yanvarev and A. L. Khandazhynskaya                            | 597–598 | <b>Base-Modified Ribonucleosides as Potential Anti-Hepatitis C virus Agents</b><br>A. V. Ivanov, O. A. Smirnova, N. A. Golubeva, M. A. Ivanov, V. L. Tunitskaya, A. V. Shipitsyn and L. A. Alexandrova                                                                                                                         | 619–620 |
| <b>Design and development of curcumin bioconjugates as antiviral agents</b><br>D. Rai, D. Yadav, J. Balzarini, E. De Clercq and R. K. Singh                                                                                                          | 599–600 | <b>Synthesis and Antiviral Evaluation of Conformationally Fixed Bicyclohexanyl Nucleosides with Ethenyl Group at C3'-Position</b><br>W. H. Kim, K. R. Kim, A.-Y. Park, H.-R. Lee, J.-A. Kang, J. Ho Bae, L. S. Jeong and H. R. Moon                                                                                            | 621–622 |
| <b>The novel unsaturated acyclic nucleoside analogues: cytostatic and antiviral activity evaluations</b><br>K. Wittine, K. Benci, S. K. Pavelić, K. Pavelić and M. Mintas                                                                            | 601–602 | <b>Synthesis of conformationally locked carbocyclic nucleoside phosphonates to probe the active site of HIV-1 RT</b><br>H. Saneyoshi, B. C. Vu, S. H. Hughes, P. L. Boyer, S. G. Sarafianos and V. E. Marquez                                                                                                                  | 623–624 |
| <b>Synthesis and study of 9-deazaguanosine derivatives as potential inhibitors of RNA virus replication</b><br>M. Hamann, C. Pierra, R. Storer, J.-P. Sommadossi, A. G. Loi, A. Cadeddu, M. Fanti, N. Boscu, F. Bassetti, M. Liuzzi and G. Gosselin  | 603–604 | <b>Ribonucleosides Bearing Michael Acceptors: Synthesis, Molecular Docking, Enzyme Inhibition Data and Antiviral Activity</b><br>V. Nair, V. Uchil, X. Ma, Q. Zhu, F. Zhang and E. Bonsu                                                                                                                                       | 625–626 |
| <b>Synthesis and antiviral activity of novel derivatives of 2'-β-C-methyleytidine</b><br>J. Fogt, P. Januszczuk, G. Framski, T. Onishi, K. Izawa, E. De Clercq, J. Neyts and J. Boryski                                                              | 605–606 | <b>Molecular mechanisms in two cell death-types, necrosis and apoptosis, induced by 5-fluoro-2'-deoxyuridine</b><br>A. Sato, A. Satake, A. Hiramoto, E. Miyazaki, A. Okamatsu, K. Nakama, O. Hiraoka, T. Miyake, H.-S. Kim and Y. Wataya                                                                                       | 627–628 |
| <b>Synthesis of Pyrimidine Analog of Fluoroneplanocin A as Potential Anti- HCV Agent</b><br>H.-R. Lee, W. H. Kim, A.-Y. Park, J.-A. Kang, P. Chun, J. H. Bae, L. S. Jeong and H. R. Moon                                                             | 607–608 | <b>Thymidine phosphorylase as a target for antiangiogenesis treatment</b><br>G. J. Peters, I. V. Bijnsdorp and M. Fukushima                                                                                                                                                                                                    | 629     |
| <b>Synthesis and antiviral evaluation of (±)-4'-ethynyl-5'-difluorocarbocyclic-d4T analogue</b><br>H. Kumamoto, K. Haraguchi, M. Ida, H. Tanaka, T. Hamasaki and M. Baba                                                                             | 609–610 | <b>A Proposal of the Structure of Modified Nucleosides Expected to be highly Anti-Viral Active and Lowly Toxic</b><br>H. Ohrui                                                                                                                                                                                                 | 631–632 |

|                                                                                                                                                                                                                                                                                                     |         |                                                                                                                                                                                                                                                                        |         |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>ppGpp analogues as antibacterial compounds</b><br>W. Ezequiel, K. Jehoshua, G. Gad and V. Roeë                                                                                                                                                                                                   | 633–634 | <b>Pyrimidine Acyclic Nucleoside Phosphonates and Phosphorylated Analogs (Part 2): Syntheses and Investigation of Their Inhibitory Effects Towards Human Thymidine Phosphorylase</b><br>K. Pomeisl, A. Holý and I. Votruba                                             | 657–658 |
| <b>Hetero-expanded Purine Nucleosides. Design, Synthesis and Preliminary Biological Activity</b><br>K. L. Seley-Radtke, Z. Zhang, O. R. Wauchope, S. C. Zimmermann, A. Ivanov and B. Korba                                                                                                          | 635–636 | <b>Synthesis of conformationally locked carbocyclic 1,3-diazepinone nucleosides as inhibitors of cytidine deaminase</b><br>O. R. Ludek, G. K. Schroeder, R. Wolfenden and V. E. Marquez                                                                                | 659–660 |
| <b>Biologically active 2-5A analogs containing 3'-O,4'-C-propylene adenosine as potent RNase L agonists</b><br>K. Morita, S. Obika, T. Imanishi, Y. Kitade and M. Koizumi                                                                                                                           | 637–638 | <b>9-Deazaguanine derivatives: synthesis and inhibitory properties as multi-substrate analogue inhibitors of mammalian PNPs</b><br>T. Yatsu, M. Hashimoto, S. Hikishima, L. Magnowska, A. Bzowska and T. Yokomatsu                                                     | 661–662 |
| <b>Synthesis and Antitumor Activity of 1-β-4-Selenoarabinofuranosyl cytosine (4'-Seleno-ara-C)</b><br>H. Lee, D. K. Tosh, W. J. Choi, Y. M. Kim, S. K. Lee and L. S. Jeong                                                                                                                          | 639–640 | <b>Thermodynamic studies of interactions of calf spleen PNP with acyclic phosphonate inhibitors</b><br>K. Breer, B. Wielgus-Kutrowska, M. Hashimoto, S. Hikishima, T. Yokomatsu, R. H. Szczepanowski, M. Bochtler, A. Girstun, K. Staroń and A. Bzowska                | 663–664 |
| <b>Design and Synthesis of Truncated 4'-Thioadenosine Derivatives as Potent and Selective A<sub>3</sub> Adenosine Receptor Antagonists</b><br>X. Hou, S. Pal, W. J. Choi, H. O. Kim, A. Tipnis, K. A. Jacobson and L. S. Jeong                                                                      | 641–642 | <b>Nucleoside Phosphonic Acids in Thymidine Phosphorylase Inhibition: Structure - Activity Relationship</b><br>N. Panova, I. Košiová, M. Petrová, V. Vaněk, R. Liboska, S. Kovačková, P. Kočalka, Š. Králíková, Z. Točík, O. Páv, O. Pačes, D. Rejman and I. Rosenberg | 665–666 |
| <b>Synthesis and Biological Activity of 7-Deaza-7-ethynyl-2'-deoxy-2'-fluoro-2'-C-methyladenosine and its 2'-C-Methyl-ribo Analogue</b><br>M. Prhac, N. Dyatkina, J. Keicher, S. Liehr, S. Koo-McCoy, D. Latour, K. Fung, K. Dunlop, J. Pouliot, T. Wang, W. Li, L. Lou, C. Roberts and R. Griffith | 643–644 | <b>Crown ether ring-fused nucleosides: synthesis and conformational properties</b><br>C. Coppola, J. D'Onofrio, G. Di Fabio, L. De Napoli and D. Montesarchio                                                                                                          | 667–668 |
| <b>Synthesis of 2-Chloro-<sup>N</sup>6-Substituted-4'-thioadenosine-5'-N, N-dialkyluronamides as Potent and Selective A<sub>3</sub> Adenosine Receptor Antagonists</b><br>W. J. Choi, H. W. Lee, X. Hou, H. O. Kim, K. A. Jacobson and L. S. Jeong                                                  | 645–646 | <b>Simple and universal method to determine dissociation constants for enzyme/ligand complexes</b><br>A. Bzowska and L. Magnowska                                                                                                                                      | 669–670 |
| <b>Effects of CpG- oligodeoxynucleotides on dendritic cell development</b><br>R. Ohue, F. Tani and N. Kitabatake                                                                                                                                                                                    | 647–648 | <b>On the analysis of fluorimetric titration curves of purine nucleoside phosphorylase</b><br>J. M. Antosiewicz, K. Breer, A. Bzowska and B. Wielgus-Kutrowska                                                                                                         | 671–672 |
| <b>Novel cell death mechanisms of trifluorothymidine</b><br>I. V. Bijnisdorp, F. Kruyt, M. Fukushima and G. J. Peters                                                                                                                                                                               | 649     | <b>Effects of Arginine Residue Introduction upon Interaction and Complexation Behavior of Peptide Ribonucleic Acids (PRNAs) with RNA: Synthesis and Properties of α-Containing Arginine</b><br>T. Wada, N. Sawa, H. Sato, S. Futaki and Y. Inoue                       | 673–674 |
| <b>In vitro activity and mechanism of action of a duplex and multidrug of ethynyleytidine and 5-fluorodeoxyuridine</b><br>I. V. Bijnisdorp, R. A. Schwendener, H. Schott, S. Schott, I. Fichtner, R. J. Honeywell, N. Losekoot, A. C. Laan and G. J. Peters                                         | 651     | <b>Photo-Switched DNA-Binding of a Photochromic Spiropyran</b><br>J. Andréasson, J. Andersson, P. Lincoln and S. Li                                                                                                                                                    | 675     |
| <b>Synthesis and Biological Evaluation of 5''-Iodoneplanocin A and Its Analogues</b><br>A.-Y. Park, K. R. Kim, H.-R. Lee, J.-A. Kang, W. H. Kim, L. S. Jeong and H. R. Moon                                                                                                                         | 653–654 | <b>Synthesis of Alkylated Poly(1-vinylimidazole) for a New pH-Sensitive DNA Carrier</b><br>T. Hakamatani, S. Asayama and H. Kawakami                                                                                                                                   | 677–678 |
| <b>Boranophosphate siRNA-aptamer chimeras for tumor-specific downregulation of cancer receptors and modulators</b><br>B. R. Shaw, L. Moussa, M. Sharaf, M. Cheek and M. Dobrikov                                                                                                                    | 655–656 | <b>Antisense Inhibition of Human Telomerase by Phosphorothioate Oligonucleotide-Peptide Conjugates</b><br>I. Diala, S. Murao and M. Fujii                                                                                                                              | 679–680 |

|                                                                                                                                                                                                                                           |         |                                                                                                                                                                                                         |         |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| <b>Single-Molecule Accommodation of Streptavidin in Nanometer-Scale Wells Formed in DNA Nanostructures</b><br>A. Kuzuya, K. Numajiri, M. Kimura and M. Komiyama                                                                           | 681–682 | <b>RNA aptamers that reversibly bind to photoresponsive peptide</b><br>G. Hayashi, M. Hagihara and K. Nakatani                                                                                          | 703–704 |
| <b>Addressable Molecular Node Assembly – High Information Density DNA Nanostructures</b><br>E. Lundberg, J. Tumpene, R. Kumar, P. Sandin, N. Gale, I. S. Nandhakumar, B. Albinsson, P. Lincoln, L. M. Wilhelmsson, T. Brown and B. Nordén | 683–684 | <b>Efficient quenching of the excimer fluorescence derived from pyrene arrays on RNA duplexes</b><br>K. Maie, M. Nakamura and K. Yamana                                                                 | 705–706 |
| <b>Self-Assembled DNA Photonic Wire</b><br>J. K. Hannestad, P. Sandin and B. Albinsson                                                                                                                                                    | 685     | <b>Zipper-like assembly of multi-pyrenes covalently attached to RNA sequences via duplex formation</b><br>M. Nakamura, Y. Murakami and K. Yamana                                                        | 707–708 |
| <b>Programmed regulation of conformational switching in diagnostic biomolecular automata using error control molecules</b><br>M. Hirabayashi, H. Ohashi and T. Kubo                                                                       | 687–688 | <b>Selective recovery of target cells employing a bi-functional aptamer</b><br>S. P. Ohuchi                                                                                                             | 709–710 |
| <b>Functionalization of DNA G-Wires for patterning and nanofabrication</b><br>S. Lyonnais, O. Piétremont, A. Chepelianski, S. Guéron, L. Lacroix, E. Le Cam and J.-L. Mergny                                                              | 689–690 | <b>Regulating mRNA translation with a kiss</b><br>E. Dausse, C. Belair, C. Staedel and J.-J. Toulmé                                                                                                     | 711–712 |
| <b>A Membrane Anchored DNA-based Energy/Electron Transfer Assembly</b><br>K. Börjesson, J. Tumpene, T. Ljungdahl, L. M. Wilhelmsson, B. Nordén, T. Brown, J. Mårtensson and B. Albinsson                                                  | 691     | <b>Improvement of transcription- and transfection-efficiency by synthetic polyampholytes</b><br>C. Yoshihara, T. Ito, C.-Y. Shew and Y. Koyama                                                          | 713–714 |
| <b>Properties of Superparamagnetic Iron Oxide Nanoparticles Assembled on Nucleic Acids</b><br>K. Tanaka, N. Kitamura and Y. Chujo                                                                                                         | 693–694 | <b>Analysis of cationic comb-type copolymers/DNA interaction by the single molecular observation and intermolecular force measurement</b><br>M. Hirano, N. Shimada, A. Kano, S. Kidoaki and A. Maruyama | 715–716 |
| <b>DX DNAs as Templates for Multiple Arrangement of Zinc Fingers</b><br>K. Sasaki, A. Onoda, M. Mizota, R. Sakamoto and T. Yamamura                                                                                                       | 695–696 | <b>Synthesis of naphthalenediimide having two <math>\beta</math>-cyclodextrins at both of its substituent termini</b><br>K. Ohtsuka, S. Sato, S. Watanabe, K. Komizo and S. Takenaka                    | 717–718 |
| <b>Light driven open/close operation of an azobenzene-modified DNA nano-pincette</b><br>X. Liang, N. Takenaka, H. Nishioka and H. Asanuma                                                                                                 | 697–698 | <b>Synthesis of new non-nucleosidic ligand building blocks for solid-phase oligonucleotide assembly</b><br>A. Jamil, E. M. Zubin and D. A. Stetsenko                                                    | 719–720 |
| <b>Preparation of coherent hetero clusters with threoninol scaffold</b><br>T. Fujii, H. Kashida and H. Asanuma                                                                                                                            | 699–700 | <b>Oligonucleotides direct synthesis on porous silicon chip</b><br>L. De Stefano, E. De Tommasi, I. Rea, L. Rotiroti, L. Giangrande, G. Oliviero, N. Borbone, A. Galeone and G. Piccialli               | 721–722 |
| <b>Incorporation of cationic dyes into DNA for distinct stabilization of duplex</b><br>H. Kashida, H. Itoh, T. Fujii and H. Asanuma                                                                                                       | 701–702 | <b>Microarray based oligonucleotide synthesis</b><br>I. A. Koslov, F. Kaper, L. Zhou and M. S. Chee                                                                                                     | 723     |
|                                                                                                                                                                                                                                           |         | <b>Erratum</b>                                                                                                                                                                                          | 725     |
|                                                                                                                                                                                                                                           |         | <b>Index</b>                                                                                                                                                                                            | 727–732 |